5,369,108

35. Young, C. W. Fanucchi, M. P., Walsh, T. B., Blatzer, L., Yaldaie, S., Stevens, Y. W., Gordon, C., Tong, W., Rifkind, R. A., and Marks, P. A. (1988) Cancer Res. 48: 7304-7309.

36. Andreeff, M., Young, C., Clarkson, B., Fetten, J., Rifkind, R. A., and Marks, P. A. (1988) *Blood* 72: 186a. 37. Marks, P. A., Breslow, R., Rifkind, R. A., Ngo, L., and Singh, R. (1989) *Proc. Natl. Acad. Sci.* (USA) 86:

38. Breslow, R., Jursic, B., Yan, Z. F., Friedman, E., 10 Leng, L., Ngo, L., Rifkind, R. A., and Marks, P. A. (1991) Proc. Natl. Acad. Sci. (USA) 88: 5542-5546.

39. Ohta, Y., Tanaka, M., Terada, M., Miller, O. J., Bank, A., Marks, P. A., and Rifkind, R. A. (1976) *Proc. Natl. Acad. Sci.* (USA) 73: 1232-1236.

40. Collins, S. J., Gallo, R. C., and Gallagher, R. E. (1978) Nature (London) 270; 405-409.

41. Synder, S. W., Egorin, M. J., Geelhaar, L. A., Hamburger, A. W., and Callery, P. S. (1988) Cancer Rex 48; 3613–3616.

What is claimed is:

6358-6362.

1. A compound having the structure:

wherein each of R_1 and R_2 are independently the same as or different from each other; when R_1 and R_2 are the same, each is a substituted or unsubstituted cycloalkylamino, pyridineamino, piperidino, 9-purine-6-amine, or thiazoleamino group; when R_1 and R_2 are different, $R_1=R_3$ —N—R4, wherein each of R_3 and R_4 are independently the same as oddifferent from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, branched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy, arylalkyloxy, or pyridine group, or R_3 and R_4 bond together to form a piperidine group and R_2 is a hydroxylamino, hydroxyl, 40 amino, alkylamino, or alkyloxy group; and n is an integer-from about 4 to about 8.

R₃-N C-(CH₂)_n-C O

wherein each of R₃ and R₄ are independently the same as or different from each other and are a hydrogen atom, a hydroxyl group, a substituted or unsubstituted, dranched or unbranched alkyl, alkenyl, cycloalkyl, aryl, alkyloxy, aryloxy, arylalkyloxy, or pyridine group, or R3 and R4 bond together to form a piperidine group; R2 is a hydroxylamino, hydroxyl, amino, alkylamino, or alkyloxy group; and n is an integer from about 4 to about 8.

3. A compound of claim 2, wherein R2 is a hydrox-

3. A compound of claim 2, wherein R_2 is a hydroxylamino, hydroxyl, amino, methylamino, or methyoxy group and in S_2 .

4. A compound of claim 3, wherein R4 is a hydrogen atom and R3 is a substituted or unsubstituted phenyl group.

15 S. A compound of claim 4, wherein the phenyl group is substituted with a methyl, cyano, nitro, trifluoromethyl, amino, aminocarbonyl, methylcyano, chloro, fluoro, promo, iodo, 2,3-difluoro, 2,4-difluoro, 2,5-difluoro, 3,4-difluoro, 3,5-difluoro, 2,6-difluoro, 1,2,3-trifluoro, 2,3,6-tertrafluoro, 2,3,4,5-tertrafluoro, 2,4-tertrafluoro, 2,3,4,5-tertrafluoro, 2,3,4,5-tert

6. A compound of claim 3, wherein R4 is a hydrogen atom and R3 is a cyclohexyl group.

7. A compound of claim 3, wherein R4 is a hydrogen atom and R3 is a methyoxy group.

8. A compound of claim 3, wherein R₃ and R₄ bond

together to form a piperidine group.

9. A compound of claim 3, wherein R₄ is a hydrogen

atom and R₃ is a wdroxyl group.

10. A compound of claim 3, wherein R₄ is a hydrogen

atom and R₃ is a benzyloxy group.

11. A compound of claim 3' wherein R₄ is a hydrogen

atom and R₃ is a 8-pyridine group.

12. A compound of claim 3, wherein R₄ is a hydrogen

The A compound of claim 3, wherein K4 is a hydrogen atom and R3 is a β-pyridine group.

13. A compound of claim 3, wherein R₄ is a hydrogen atom and R₃ is a α -pyridine group.

14. A compound of claim 3, wherein R₃ and R₄ are both methyl groups.

15. A compound of claim 3, wherein R₄ is a methyl group and R₃ is a phenyl group.

16. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of the compound of claim 2.

17. A pharmaceutical composition of claim 16 in combination with an antitumor agent.

Badd

60

(<u>20</u>)